

Periodic function, trip 3 (easy)

Solar cycle predictions are used by various agencies and many industry groups. The solar cycle is important for determining the lifetime of satellites in low-Earth orbit, as the drag on the satellites correlates with the solar cycle [...]. [\(NOAA\)](#)

[\(Solar Cycle\)](#)

Sunspot Number Progression : Observed data through May 2008 ; Dec 2012 ; Nov 2014

The goal of the problem is to propose a perfect prediction center, with weak constraints.

Let us consider periodic functions from \mathbf{Z} to \mathbf{R} .

```
def f(x): return [4, -6, 7][x%3] # (with Python notations)
# 4, -6, 7, 4, -6, 7, 4, -6, 7, 4, -6, 7, 4, -6, 7, ...
```

For example, f is a 3-periodic function, with $f(0) = f(3) = f(6) = f(9) = 4$.

With a simplified notation we will denote f as $[4, -6, 7]$.

For two periodic functions (with integral period), the quotient of periods will be rational, in that case it can be shown that the sum of the functions is also a periodic function. Thus, the set of all such functions is a vector space over \mathbf{R} .

For that problem, we consider a function that is the sum of several periodic functions all with as period an integer N at maximum. You will be given some starting values, you'll have to find new ones.

Input

The first line contains an integer T , the number of test cases, then each case will be given on three lines.

On the first line, you will be given an integer N .

On the second line, you will be given integers y : the first (0-indexed) $N \times N$ values of a periodic function f that is sum of periodic functions all with as period an integer N at maximum.

On the third line, you will be given $N \times N$ integers x .

Output

Print $f(x)$ for all required x . See sample for details.

Example

Input:

```
2
2
5 7 5 7
6 7 8 9
3
15 3 17 2 16 4 15 3 17
```

10 100 1000 10000 100000 1000000 10000000 100000000 1000000000

Output:

5 7 5 7

16 16 16 16 16 16 16 16 16

Explanation

Test case 1: for example f can be seen as the sum of two periodic functions : $[5] + [0, 2]$ (with simplified notations)

We know that $f(0)=5$ and $f(1)=7$, we can deduce that $f(6)=5$, and so on...

Test case 2: for example f can be seen as the sum of three periodic functions : $[10] + [5, -8] + [0, 1, 2]$ (with simplified notations). In that case $f(10) = [10][10\%1] + [5, -8][10\%2] + [0, 1, 2][10\%3] = 10 + 5 + 1 = 16$, and so on.

Constraints

$0 < T < 1024$

$1 < N < 14$: uniform distribution

$\text{abs}(y) < 10^9$

$0 < x < 10^9$

Information

Constraints allow easy coding with various languages. (Edited 2017-02-11 ; with compiler changes)

There's two input files, a small one and a bigger.

My PY3.4 code ended in $0.02+0.28 = 0.30$ s. My C code in 0.01s.

If you find the constraints too weak, please consider [PERIOD3](#). **Have fun ;-)**